

About famous blindsight patient FS

Among the three patients in our study (Elliot et al., 2014) (RP, FS, and LE), one is the well-known patient FS, born in 1937, who has become famous in particular through the work of Stoerig and colleagues. Quoting Stoerig et al. (1998), “FS is one of the most extensively studied blindsight patients (see Stoerig & Cowey, 1997, for review).” Here are a few references as pointers.

References

- Elliott, M. A., Seifert, D., Poppel, D. A. & Strasburger, H. (2015). Transient increase of intact visual field size by high-frequency narrow-band stimulation. *Consciousness and Cognition* 32, Special Issue: Type 2 Blindsight, 45–55.
- Kleiser, R., Wittsack, J., Niedeggen, M., Goebel, R. & Stoerig, P. (2001). Is V1 necessary for conscious vision in areas of relative cortical blindness? *NeuroImage*, 13, 654–661.
- Stoerig, P. (1987). Chromaticity and achromaticity. Evidence for a functional differentiation in visual field defects. *Brain* 110, 869–886.
- Stoerig, P. (1993). Spatial summation in blindsight. *Visual Neuroscience* 10, 1141–1149.
- Stoerig, P. & Cowey, A. (1997). Blindsight in man and monkey. *Brain*, 120, 535–559.
- Stoerig, P., Kleinschmidt, A. & Frahm, J. (1998). No visual responses in denervated V1: high-resolution functional magnetic resonance imaging of a blindsight patient. *NeuroReport* 9, 21–25.
- Goebel, R., Muckli, L., Zanella, F. E., Singer, W. & Stoerig, P. (2001). Sustained extrastriate cortical activation without visual awareness revealed by fMRI studies of hemianopic patients. *Vision Research* 41, 1459–1474.